FEDOROV, V.1

PHASE I BOOK EXPLOITATION

807/4103


- Levin, Moisey Yevseyevich, Georgiy Andreyevich Malinin, Mikhail Mikolayevich Mandrazhitskiy, Valentin Petrovich Sinitsyn, and Valeriy Ivanovich Fedorov
- Zashchita it sredstv massovogo porazheniya (Protection Against Means of Mass Destruction) 2nd ad, Moscow, Uchpedgiz, 1960. 176 p. 50,000 copies printed.
- General Ed.: V. P. Sinitsyn, Candidate of Technical Sciences, and G. A. Malinin. Ed.: A. A. Korotkiy; Tech. Ed.: R. V. Tsyppo.
- PURPOSE: This book is intended for the public instructors of PVO DOSAAF (Air Defence Organization under the All-Union Voluntary Society for the Promotion of the Army, Aviation and Navy).
- COVERAGE: The book gives fundamental information on atomic, chemical, and bacteriological weapons and on means of individual and collective protection. It states that the problem has been studied sufficiently and that at the present time adequate means of protection exist for a well-organized and trained population. No personalities are mentioned. There are no references.

Card 1/3


			4	
Pro	ect:	ion Against Means of Mass Destruction SOV/4103		
		CONTENTS:		
Int	ogno	stion		
Ch.	ı.	Modern Means of Attack From the Air	5	:
Ch.		Atomic Weapons and Their Casualty Effect	5 15	
Ch.		High-Explosive, Fragmentation, and Incendiary Weapons	41	
Ch.		Chemical Weapons and Their Casualty Rffect	51	
Ch.		Bacteriological Weapons and Their Casualty Effect	59	I
		Protective Equipment for the Individual	71	
		Protective Equipment for Groups	86	
Ch.	8.	Protection of Food, Water, and Forage Against Contamination by Poisonous, Radioactive, or Bacteriological Substances	99	
Card	2/3		77	

Protecti	ion Against Means of Mass Destruction 807/4103	
Ch. 9.	Tasks and Organization of the Local Air Defense Relative to Dwellings, Establishments, Institutions, and State and Collect Farms. Rules of Conduct and Action for the Population According to the Signals of the Local Air Defense	ive ing
Ch. 10.	Reconnaissance of Stricken Areas	110
Ch. 11.	Emergency and Rescuing Operations in Stricken Areas	130
Ch. 12.	Fire Prevention Measures; Extinguishing Fires in Progress and Breaking Out	136
ch, 13.	Methods and Means of Decontamination; Degassing and Disinfect	ion 145
Ch. 14.	Duties of Personnel of Salf-Defense Groups Responding to Signa Given by the Local Air Dufense	ls 166
AVAILABL	E: Library of Congress (UA926.138 1960)	
Card 3/	- A	C/rn/ec -25-60

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7



"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7



SOV /137-58-12-25247

Translation from: Referativnyy zhurnal. Metallurgiya, 1958, Nr 12, p 172 (USSR)

a Kand. Tekhn. nauh.

AUTHORS: Fedorov, V. K. , Zhavoronkova, R.S.

TITLE:

Improvement of Casting Properties of Kh28 Alloy (Uluchsheniye

liteynykh svoystv splava Kh28)

PERIODICAL: Sb. statey Vses. n.-i. i konstrukt. in-t khim. mashinostr., 1957, Vol 23, pp 3-13

ABSTRACT: An investigation was made of heat tearing and "black-spot" formation in castings of Kh28 alloy of the following composition (in %): C 0.5-1.0, Mn 0.5 - 0.8, Si 0.5 - 1.3, and Cr 26 - 30, depending upon the structure of the casting, pouring temperature, melting procedure, composition of the charge, conditions of inoculation, and rate of cooling. The following specimens were cast: 25x25 mm in cross section, bent at 120, 90, and 30-degree angles, specimens 30 and 60 mm in diam and a rake-shaped specimen, as well as standard specimens for bending tests and for determination of fluidity. The following inoculants were used: 75% FeSi in amounts of 0.5 and 1% of the weight of the metal and a mixture of FeSi+FeTi (1:1) 0.6 - 0.8% of the weight of the metal. The character of the fracture, the microstructure, and

Card 1/2

SOV /137-58-12-25247

Improvement of Casting Properties of Kh28 Alloy

the mechanical properties were investigated. It was established that to avoid hot tearing and "black spots" higher casting temperature is necessary: >1500°C for thin-wall and >1450° for thick-wall castings. In order to produce finer grain and to improve the mechanical properties the authors recommend inoculation with the FeS+FeT1 (1:1) mixture in amounts of 0.6-0.8% of the weight of metal, and an increase of the cooling rate by chill casting and setting up a cooling system.

 $\mathbf{T} \cdot \mathbf{F}$.

Card 2/2

Casting of ring pots and heat stabilizing treatment of compressor piston rings. Sbor.st.MIRHIMMASH no.23:47-58 '57.

(Molding (Founding)) (Piston rings)

FEDOROV, V.K.

by Phase I book exploitation 30V/5488

- Moscow. Vsesoyuznyy nauchno-issledovatel'skiy i konstruktorskiy institut khimicheskogo mashinostroyeniya.
- Materialy v khimicheskom mashinostroyenii (Materials in Chemical Machine Building) Moscow, Informatsionno-izdatel'skiy otdel, 1960. 143 p. (Series: Its: Trudy, vyp. 34) 3,000 copies printed.
- Sponsoring Agency: Gosudarstvennyy komitet Soveta Ministrov SSSR po avtomatizatsii i mashinostroyeniyu and Vsesoyuznyy nauchno-issledovatel'skiy i konstruktorskiy institut khimicheskogo mashinostroyeniya NIIKhIMMASh.
- Ed. (Title page): V. K. Fedorov, Candidate of Technical Sciences; Editorial Council: Chairman: V. B. Nikolayev; Deputy Chairman: Yu. M. Vinogradov, Candidate of Technical Sciences; B. N. Borisoglebskiy, A. N. Goncharov, Yu. G. Popandopulo, I. N. Yukalov, Candidate of Technical Sciences, and G. M. Yusova, Candidate of Technical Sciences; Ed.: V. I. Glukhov; Tech. Ed.: P. A. Vshivtsev.

Card 1/5

Materials in Chemical (Cont.)

SOV/5488

PURPOSE: This collection of articles is intended for technical personnel in chemical machine building and other branches of the machine and instrument industry.

COVERAGE: The collection deals with the results of investigations on the mechanical, corrosive, and engineering qualities of certain alloys. Also discussed are heat-treatment regimes, the phase composition of stainless steels, methods of checking products, and new designs of apparatus used in checking. References accompany each article.

TABLE OF CONTENTS:

Gavrilov, V. M. [Engineer], and <u>V. K. Fedorov</u> [Candidate of Technical Sciences]. Crystallization of Alloys in the Elastic-Vibration Field

Moskvin, N. I. [Engineer]. Metal Which Will Resist Corrosion in Molten Type Metal Containing Zinc

12

Card 2/5

Materials in Chemical (Cont.)	sov/5488	
Shapiro, M. B. [Engineer], and V. M. I Hardening of Small-Module Pinions of	Makarov [Engineer]. Induct [Speed] Reducers	1on 26
Chernykh, N. P. [Engineer, Irkutskiy in Irkutsk branch of NIIKhIMMASH]. Invertigation of Certain in Molchanova and M. I. Mil' took part in	stigation of the Effect of Steels (Engineers V. D.	33
Akshentseva, A. P. [Candidate of Techi Shumratova [Engineer]. Effect of Heat Composition of 1Kh18N9T and Kh18N12M3 P. T. Dmitriyev, B. N. Shevelkin, A. I and L. Ye. Lobanova took part in the	t Treatment on the Phase I Steels [V. N. Dayatlova, M. Shabanova, Z. K. Ogurtso	va, 50
Dyatlova, V. N. [Engineer], and le. M. pendence of the Corrosion Resistance of Steels on the a-Phase Content	. Frolikova [Engineer]. De- of 1Kh18N9T and Kh18N12M3T	- 69
Shevelkin, B. N. [Candidate of Technic Various a-Phase Contents in 1Kh18N9T S Card 3/5	eal Sciences]. Effect of Steel and α -Phase	

Materials in Chemical (Cont.)

SOV/5488

Contents in Kh18N12M3T Steel on Their Formability [Engineers A. P. Golovanova, L. L. Kravchenko, V. N. Dyatlova, and Candidate of Technical Sciences A. P. Akshentseva took part in the investigation]

82

Khakhlova, N. V. [Junior Scientific Worker], N. S. Dombrovskaya [Doctor of Chemical Sciences], V. G. Kuznetsov [Doctor of Chemical Sciences], and Ye. M. Zhilina [Engineer]. Chemical Investigation of the a-Phase Precipitated From 1Kh18N9T Steel [X-ray phase analysis was carried out at the Institute of General and Inorganic Chemistry of the Academy of Sciences of the USSR by V. G. Kuznetsov and Z. V. Popova]

Yesilevskiy, V. P. [Engineer], and N. S. Akulov [Academician of the Academy of Sciences of the Belorussian SSR]. Ponderomotive Magnetic Method of Determining the a-Phase Content in Austenitic Steel [Equipment was manufactured by NIIKhIMMASh; Technician V. M. Malinin participated in working out the electrical circuit for the a-phasometer]

Card 4/5

Materials in Chemical (Cont.)

SOV/5488

Khimchenko, N. V. [Candidate of Technical Sciences], and V. N. Prikhod'ko [Engineer]. Wide-Range Ultrasonic Analyzer for Checking the Structure of Metals [Technicians V. N. Maragayev and N. N. Materanskiy participated in the production of the attachment] 120

Khimchenko, N. V. and V. N. Prikhod'ko. Use of the Wide-Range Ultrasonic Analyzer in Investigating the Structure of Steel and Cast Iron

130


Khimchenko, N. V., V. N. Prikhod'ko, and V. P. Gozak [Engineer]. Checking the Metal Quality of Large Shafts Under Factory Conditions

137

AVAILABLE: Library of Congress

VK/wrq/jw 9-26-61

Card 5/5



62121

S/170/61/004/006/013/015 B129/B212

11.7430

AUTHORS:

Leont'yev, A. I., Fedorov, V. K.

TITLE:

Calculation of the one-dimensional flow of a gas in a cylindrical channel for a given law of the heat supply

PERIODICAL:

Inzhenerno-fizicheskiy zhurnal, v. 4, no. 6, 1961, 125-127

TEXT: A solution is given for the problem of the one-dimensional flow of a compressed gas in a cylindrical channel for the case where the coefficient of the hydraulic resistance is constant along the pipe. If strong heat flows and great velocities of the gas flow occur it is necessary to take into account the effect of the temperature factor and the number M on the coefficient of resistance. The authors compare graphically their calculation results with those of other researchers. It is shown that consideration of the compression effect on the coefficient of friction at supersonic velocities of the gas flow will essentially affect the law describing the change of λ ($\lambda = \omega/a^2 = \text{velocity}$ of the gas flow; viz. critical velocity) along the pipe. The divergence

Card 1/4

23757

Calculation of the one-dimensional...

S/170/61/004/006/013/015 B129/B212

of λ will increase if M increases at the entrance. The maximum pipe lengths at supersonic speeds of the gas at the entrance of the pipe are comparatively short, and the problem of the expansion of the one-dimensional flow diagram needs further studies for these conditions.

$$\xi = \xi_0 \left(1 - \frac{k-1}{k+1} \lambda^2 \right)^{0.6} \sqrt{\frac{T_{cr}}{T_0}}, \tag{1}$$

From the results shown in Fig. 2 it is apparent that the effect of the compressibility on the coefficient of friction is given by the change of the critical pipe length for supersonic speeds. Fig. 2 shows the critical length of the pipe as a function of the reduced velocity at the entrance. The dotted curve is taken from S. A. Khristianovich (Prikladnaya gazovaya dinamika (Applied Gas Dynamics), 1948).

S. S. Kutateladze and F. S. Voronin are mentioned. There are 3 figures and 4 Soviet-bloc references.

Card 2/4

:23757 \$/170/61/004/006/013/015 B129/B212

Calculation of the one-dimensional...


ASSOCIATION: Energeticheskiy institut im. G. M. Krzhizhanovskogo, g. Moskva (Institute of Power Engineering imeni

G. M. Krzhizhanovskiy, Moscow)

SUBMITTED:


September 22, 1960

λ as function of Fig. 2: the pipe length calculated by the author (6) and by Khristianovich (A).



Card 3/4

CIA-RDP86-00513R000412630008-7" APPROVED FOR RELEASE: 03/20/2001



25556 S/170/61/004/008/006/016 B116/3201

26.5200

AUTHORS:

Leont'yev. A. I., Fedorov, V. K.

TITLE:

Effect of inlet conditions upon the law of heat exchange in the initial section of a cylindrical tube

PERIODICAL:

Inzhenerno-fizicheskiy zhurnal, v. 4, no. 8, 1961, 63 - 68

TEXT: The results of an analysis of experimental data of VTI, MEI, and ENIN concerning convective heat transfer in the initial section of a cylindrical tube are presented. The analysis was made on the basis of local simulation. The heat-exchange laws for various conditions at the tube inlet were established. Methods of calculating the convective heat exchange in the initial section of the cylindrical tube are presented for the case $T_{\rm CT}$ = const and $q_{\rm CT}$ = const. The fundamental ideas of the theory of local simulation have been presented in papers by V. M. Ievlev (Refs. 1 and 2: DAN SSSR, t. 36, no. 6, 1952 and DAN SSSR, t. 37, no. 1, 1952). The equation of the thermal boundary layer for the initial section of a cylindrical tube reads:

Card 1/10

(2)

25556 S/170/61/004/008/006/016 B116/3201

Effect of inlet conditions ..

$$\frac{dRe_0}{dRe_x} + Re_0 \frac{1}{\overline{l_0}} \frac{d\overline{l_0}}{dRe_x} = \alpha_m,$$

$$Re_{\theta} = \frac{\overline{u}}{\overline{\mu_{\theta}}} \int_{0}^{\Delta} \frac{\rho u}{\overline{u}} \left(1 - \frac{t_{\theta}}{\overline{t_{\theta}}} \right) \left(1 - \frac{y}{r} \right) dy;$$

where $\overline{l_0} = \overline{T_0} - T_{cr}$; $l_0 = T_0 - T_{cr}$;

$$dRe_x = \frac{\rho u dx}{\mu}; \quad \alpha_m = \frac{q_{cr}}{g \rho u c f_0}$$

 T_o and T_o denote the temperature found when decelerating the gas in the flow center and in the boundary layer, respectively. \overline{Q} and $\overline{\mu}$ are, respectively, the density and viscosity with respect to the thermodynamic Card 2/10

25556 S/170/61/004/008/006/016 B116/B201

Effect of inlet conditions

temperature in the flow center. u is the velocity in the undisturbed flow. r and x denote the tube radius and the distance from the tube txis, respectively. For solving (1), it is necessary to determine the relationship between α_m and Re_{θ} . If, during the experiments, the distribution of the specific heat flows, of the wall temperature, and of the static pressures along the tube are measured, the local values of

 Re_0 and α_m can be determined on the strength of these measurements and

from the following formulas:

$$Re_{\theta} = \frac{\int_{0}^{\pi} q_{cr} dx}{\overline{\mu} c_{\rho} g t_{\theta}},$$

$$\alpha_{m} = \frac{q_{cr} D}{Re_{D} c_{\rho} g \overline{\mu} t_{\theta}},$$
(4)

Card 3/10

25556 B/170/61/004/008/006/016 B116/B201

Effect of inlet conditions ...


$$Re_{D} = \beta (1 - \beta^{1})^{\frac{1}{h-1}} Re_{D_{0}};$$

$$\beta = \frac{\overline{u}}{w_{\text{max}}}; \quad w_{\text{max}} = \sqrt{2c_{\rho}T_{0}}; \quad Re_{D_{0}} = \frac{\rho_{0} w_{\text{max}} D}{\rho_{0}}$$

where μ_0 is the dynamic viscosity with respect to the impact temperature. The value of the dimensionless velocity β is determined on the basis of the distribution of static pressures and from the relation

$$p/p_0 = (1 - \beta^2)^{k/(k+1)}$$
 (6)

As may be seen from Fig. 1, conditions at the tube inlet have a considerable effect upon the heat exchange in the initial tube section. The equation for the thermal boundary layer, the equation of continuity, and the law of heat exchange are used to derive the calculation formulas. For the case $T_{\rm CT}$ = const, one obtains the formula



Effect of in	alet conditions	955 8/170/61/ B116/B201	556 004/008/006/016	X
and for the	case q _{CT} = const the for	rmula	•	
	, and the second	 In the process of the contract of	The second of	
<u> </u>	$= \frac{1}{2 \cdot 5.2 \frac{\text{Nu}_1}{\text{Prof.4}} A} \left[-(5.2 \text{ Pr}^{0.6}) \right]$	$ARe_{\theta} + ARe_{D_1} - 0.5 \frac{Nu_1}{Pr}$	- Re	
	$2 \cdot 5.2 \frac{101}{Pr^{0.4}} A$	••	(18)	
	/ ¥	Nu12 Nu.		45
+1/	$\int 5.2 \text{Pr}^{0.6} A \text{Re}_{0} + A \text{Re}_{D_{0}} - 0.5$	$\frac{Re_{\bullet}^{m}}{Pr}$ Re $\frac{Al}{Pr^{4.4}}$ Al	₹e#***].	
. ,		O ST. for case III	Fig. 1.	
	, Fig. 1, A = 0.214, m = m = 0.32. When derivin			he 55
	A L	'ANBREH WALL CALELL SI	TAO MODOMMA -5	
means of a	formula (not given here	mba mathad presen	nted here may al	
	L. IL. ASSA OF SMY IDW O	IONCHENIUM MODE BWD	NET COLOR	be.
		I THE DIGDIEM OF A	NA ATTACA	et
conditions	upon convective heat exc	change does not see	n to be delinite	73
Card 6/10		•		
				.5.

.10

13

25556 S/170/61/004/008/006/016 B116/B201

Effect of inlet conditions ...

solved, considering that only three different cases were examined. There are 3 figures and 6 Soviet-bloc references.

ASSOCIATION: Energeticheskiy institut im. G. M. Krzhizhanovskogo, g. Moskva (Institute of Power Engineering imeni G. M. Krzhizhanovskiy, Moscow)

SUBMITTED: October 21, 1960

Fig. 1: Heat-exchange laws established on the basis of experimental data. Legend: (I) V. V. Kirillov (Ref. 4: Kandidatskaya dissertatsiya. MEI, 1958) (MEI); (II) V. L. Lel'chuk and B. V. Dedyakin (Ref. 3: Voprosy teploobmena. Izd. AN SSSR, 1959); (III) ENIN (Ref. 5: Kalikhman L. Ye. Turbulentnyy pogranichnyy sloy na krivolineynoy poverkhnosti, obtekayemoy gazom. Oborongiz, 1956); (TV) data concerning a plate (am

card 7/10

37561

26.5200

5/096/62/000/005/008/009 E194/E454

AUTHORS:

Kosterin, S.I., Doctor of Technical Sciences, Professor,

Leont'yev, A.I., Candidate of Technical Sciences,

Federov, V.K., Engineer

TITLE:

Methods of generalizing experimental data on

convective heat transfer during motion of gas in the

initial section of a tube

PERIODICAL: Teploenergetika, no.5, 1962, 70-72

A review of existing methods of generalizing experimental data on heat transfer by convection with a turbulent boundary % layer which are based on criterial equations shows that none of them is reliable or scientifically well-founded. accordingly recommended to use the theory of local modelling, according to which the object of the experiment is to establish the laws of heat transfer and resistance in the turbulent boundary layer; the influence of various external factors such as pressure distribution and wall temperature being allowed for in the boundary layer equations. Equations of the thermal boundary layer for the motion of gas in the initial section of a tube are Card 1/4

S/096/62/000/005/008/009 E194/E454

Methods of generalizing ...

then written down. To obtain local experimental values of Stanton's criterion it is necessary to determine the gas parameters in the body of the flow, which may be done either from measurements of static pressure distribution over the length of the tube or from thermal measurements alone. The derivation of the following expression for the Stanton and Pekle criteria is explained

$$Pe_{\theta} = \frac{\int_{0}^{x} q_{ct} dx}{t_{o} \lambda_{oo}}$$
 (2)

$$St = \frac{q_{ctD}}{(Re_{D_1} + 4h Re_{v}) Pr \lambda_{oo}^{t}}$$
 (6)

where q_{ct} - heat flow at the tube wall; $t_0 = T_{ct}^H - T_{ct} = equilibrium wall temperature - wall temperature;$ Card 2/4

Methods of generalizing ...

S/096/62/000/005/008/009 E194/E454

 λ_{00} - coefficient of thermal conductivity at the retardation temperature $T_{00};\ D$ - tube diameter. The suffix θ relates to the width of impulse loss. These two expressions may be used to determine local values of the Pekle and Stanton criteria and to establish the law of heat transfer. It is found that all available experimental data obtained in several different ways lie around a single curve corresponding to the equation

$$St = \frac{0.014(1 - \beta^2)^{0.5}}{Pe_{\Theta}^{0.25} \overline{T}_{ct}^{0.5} Pr^{0.5}}$$
(7)

where

$$\beta = \frac{u_0}{\sqrt{2c_pT_{00}}}$$

 u_0 - initial gas speed. This method of generalising experimental data has the advantage over criterial treatment because it is possible to exclude the influence of the criteria x (which Card 3/4

Methods of generalizing ...

S/096/62/000/005/008/009 E194/E454

mainly governs the distribution of heat transfer coefficients over the length of the tube) on the laws of heat transfer and the influences of temperature variations and compressibility can be expressed directly. The proposed law of heat transfer is of universal nature and the direct influence of x and of the law of application of heat in the distribution of local heat transfer coefficients is allowed for by the equation of the thermal boundary layer. There is I figure.

ASSOCIATIONS: Institut mekhaniki AN SSSR (Institute of Mechanics AS USSR)

Institut teplofiziki SO AN SSSR (Institute of

Thermal Physics SO AS USSR)

Card 4/4

ALEKSEYEVA, M.S.; FEDOROV, V.K.

Results of the study of higher nervous activity in three generations of rats, the progenitors of which were irradiated with Co 60 gamma rays. Med. rad. 8 no.3:50-57 Mr '63. (MIRA 17:9)

Is Instituta fiziologii imeni Pavlova AN SSSR.

ACCESSION NR: AR4018336

8/0:137/64/000/001/1084/1084

SOURCE: RZh. Metallurgiya, Abs. 11538

AUTHOR: Frolov, N. A.; Belinkiy, A. L.; Fedorov, V. K.; Istrina, Z. F.

TITLE: The properties of new foundry corresion-resistant (stainless) steel, type Kh17M2TL and the area of its application in chemical machine building

CITED SOURCE: Tr. Vses. n.-1. 1 konstrukt. in-t khim. mashinostr., vy*p. 43, 84-87

TOPIC TAGS: stainless steel, corrosion resistant steel, corrosion resistant steel

TRANSLATION: Steel has higher casting properties than Cr-Ni-steel of the austenitic class. Casting shrinkage determined on an instrument designed by Bol'shakov amounts to 2.12-2.21%. The flowability was determined according to a spiral probe (with a pouring temperature of 1,400 degrees the length of the spiral is equal to 300 mm; at 1,600 degrees, it is equal to 740 mm). The internal shrinkage blisters were studied on conical and cylindrical probes. In the former, a concentration of shrinkage blisters forms; in the latter, there is a large zone of shrinkage porosity, Card 1/2

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7

. gr %r	, to prove the second	
ACCESSION NR: AR4018336		
780 degrees for 2 hours) of should be conducted for the	steel does not influence it	s mechanical properties and
Dropensity for intermedati	THE CABUTHE	Buresses and for averting
in 74% boiling acetic acid a is recommended as a substitu	nd at 78% thermic phosphuri	c acid at 100 degrees, and
is recommended as a substitu	te for Or-Ni-Steel type 18-	8.
	SUB CODE: NA	Riverno 00
ر الشريخ المراجع المستقبل المراجع المستحد الم		
	Brokelin John Shi	
Card 2/2		
The second secon	men man a sala alka alam a a a a a a a a	
		The second secon

KUTATELADZE, S.S.; LEONT'YEV, A.I.; RUETSOV, N.A.; GOL'DSHTIK,
M.A.; VOLCHKOV, E.P.; DAVYDOVA, M.V.; DRUZHININ, S.A.;
KIRILLOVA, N.N.; MALENKOV, I.G.; MOSKVICHEVA, V.N.;
MIRONOV, B.P.; MUKHIN, V.A.; MUKHINA, H.V.; REEROV, A.K.;
FEDOROV, V.K.; KHABAKHPASHEVA, Ye.M.; SHTOKOLOV, L.S.;
SHPAKOVSKAYA, L.I., red.

[Heat and mass transfer and friction in a turbulent
boundary layer] Teplomassoobmen i tronie v turbulentnom
pogranichnom sloc. Novosibirsk, Red.-izd. otdel Sibirskogo otd-niia AN SSSR, 1964. 206 p. (MIRA 18:1)

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7

ACCESSION NZ: AP5006227 Pd-1/Pr-4/PG-4/F1-L W: S/3170/65/008/002/0198/0203	
73 1	
TITIE: An engineering method for calculating convective news	
courge: Inzhenerno-fizichaskiy zhurnal, v. 8, no. 2, 1955, 198-203	
The race gas flow, heat transfer, convective under, flow around sphere	
to engineering method for calculating	
distribution of the local heat transfer coefficient of the local heat transfer coefficient of the local heat transfer coefficient $\frac{3A\bar{x}(T_{ir_i}-1+A\bar{x})}{(T_{cr_i}-1+A\bar{x})^2-(T_{cr_i}-1)^2}$.	
$Nu_L = 0.47 \sqrt{Re_{L_1}} \sqrt[3]{Pr} \bar{x} \left[\left(1 - \sqrt{1 - \bar{x}^2}\right) \left(1 - \bar{x}^2\right) \right]^{-\frac{1}{2}}$	
Cord 1/2	

"APPROVED FOR RELEASE: 03/20/2001 CIA-

CIA-RDP86-00513R000412630008-7

ACCESSION NR: AP5006227

e transverse flow around a cylinder

Nu_D = 0.664 V/Re_D V/Pr sin²⁷

and at the frontal point of a cylinder

Nu_D = 0.94 V/Re_D V/Pr rec Re_D = \frac{\rho}{\rho} \frac{\rho}{\rho

Card 1/3


09010081

EWT(d)/EWT(1)/EWP(m)/EWT(m)/EWP(w)/EPF(c)/ETC/EPF(n)=2/EWG(EWA(d)/T-2/EWP(k)/FCS(k)/EWA(h)/ETC(m)/EWA(1)ACCESSION NR: AP5020939 UR/0170/65/009/002/0171/0176 536.244 63 AUTHOR: Fedorov. V. K. ; Litinskiy, E. M. W.67 TITLE: Investigation of the heat transfer of a rectangular wedge in transverse gas flow 21.44.95 SOURCE: Inzhenerno-fizicheskiy zhurnal, v. 9, no. 2, 1965, 171-176 TOPIC TAGS: convective heat transfer, heat transfer coefficient, gas flow, transverse flow, flow angle, angle of attack, wedge body, turbulent boundary layer, boundary layer flow ABSTRACT: Most of published studies on the heat transfer of complex-shape solids with forced convection are devoted to the determination of the mean value of the heat transfer coefficient. In order to obtain a correct solution in certain scientific and technological problems, however, it is necessary to know the distribution of the local heat transfer coefficients. Studies on the heat exchange of a wedgerin transverse flow have been insufficient. The Eckert solution (Eckert, E. VDE -- Forschungsheft, 416, 1942) holds

APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7"

true only for a 45° angle-of-attack. The present author uses the theory of local simulation

2 nana ((
L 5151-66	t a couldness of the first material areas and asset of the second consequence of the second cons	grania in a summa din servici Managemin e	
ACCESSION NR: AP50209	39)
and the formulas			
and the formulas	A second	(0)	
	$Pe_{\theta} = \int q_{a,\tau} dx/(T_{\theta,\tau} - T_{\theta}) \wedge_{\theta},$	(2)	
	$Peo = \int_{\mathbb{R}} q_{m,q} dx / (T_{m,q} - T_{o}) \lambda_{o},$ $St = e/\rho_{o} u_{o} c_{g} g.$	(3)	
4	ar = athense to	, (0)	
to analyze the experimenta	d data dealing with angles-of-attack from 90	o to less than 45°	
	> 4 >45 angles-of-attack, the heat trans		
Results show that with 90°			
Results show that with 90°			
Results show that with 90° expressed by	> \(> 45\) angles-of-attack, the heat trans St =0,36/Pet Pr'/.	sfer law may be	
Results show that with 90° expressed by Gas flow is then investigate	> 9 > 45° angles - of - attack, the heat trans St = 0,36/Pet Pr'/. ed with uniform distribution of temperatures	(4) and flow rates in—	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo	> 9 > 45° angles-of-attack, the heat trans St = 0,36/Per Pr". ed with uniform distribution of temperatures tion in the incoming flow is considered isoe	(4) and flow rates in tropic with the	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo geometric aspects being deshow satisfactory agreeme	St =0,36/Pe ₁ Pr". ed with uniform distribution of temperatures tion in the incoming flow is considered isoestermined by the geometry of the wedge. That with the experimental data. Experimental	(4) and flow rates in tropic with the theoretical data are also	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo geometric aspects being do show satisfactory agreement obtained for heat transfer.	St =0,36/Per Pr//. ed with uniform distribution of temperatures tion in the incoming flow is considered isoes termined by the geometry of the wedge. The not with the experimental data. Experimental data of =0.0° angle=of=attack, when turbulent	(4) s and flow rates in ntropic with the ne theoretical data aldata are also	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo geometric aspects being deshow satisfactory agreeme obtained for heat transfer aboundary of the wedge. The	St =0,36/Pe, Pr/. ed with uniform distribution of temperatures tion in the incoming flow is considered isoestermined by the geometry of the wedge. That with the experimental data. Experimental at a q=0.0° angle-of-attack, when turbulent is experimental setup and the procedure are	(4) s and flow rates in ntropic with the ne theoretical data aldata are also	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo geometric aspects being do show satisfactory agreement obtained for heat transfer.	St =0,36/Pe, Pr/. ed with uniform distribution of temperatures tion in the incoming flow is considered isoestermined by the geometry of the wedge. That with the experimental data. Experimental at a q=0.0° angle-of-attack, when turbulent is experimental setup and the procedure are	(4) s and flow rates in ntropic with the ne theoretical data aldata are also	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo geometric aspects being do show satisfactory agreeme obtained for heat transfer aboundary of the wedge. The art. has: 4 figures and 15	St =0,36/Pe, Pr/. ed with uniform distribution of temperatures tion in the incoming flow is considered isoestermined by the geometry of the wedge. That with the experimental data. Experimental at a q=0.0° angle-of-attack, when turbulent is experimental setup and the procedure are	(4) s and flow rates in ntropic with the ne theoretical data aldata are also	
Results show that with 90° expressed by Gas flow is then investigate an incoming flow. Gas mo geometric aspects being deshow satisfactory agreeme obtained for heat transfer aboundary of the wedge. The	St =0,36/Pe, Pr/. ed with uniform distribution of temperatures tion in the incoming flow is considered isoestermined by the geometry of the wedge. That with the experimental data. Experimental at a q=0.0° angle-of-attack, when turbulent is experimental setup and the procedure are	(4) s and flow rates in ntropic with the ne theoretical data aldata are also	



TITLE: Properties of tan	the summer of Y	9.
SOURCE: Ref. zh. Machino machin. Gidropriovi, Abs	costroitel'nyye materialy, konstruktsii i raschet detaley	
EF SOURCE: Tr. Vses. n.	ai. i konstrukt. in-t khim. mashinostr., vyp. 47, 1964, 39.	-49
ABSTRACT: Literature dat	its on the physical and corresional properties of its and no	0
From an investigation of 3000 are presented, as a rolling, tube expanding, of brand TH3 and on Nb.	The and Ta mechanical properties at temperatures from -70 to are data on the influence of technical operations (bending, stamping, and welding) on the corrosional properties on Ta 3 illustrations, 5 tables. /Translation of abstract/	a ii a angle
from an investigation of +300C are presented, as a	are data on the influence of technical operations (bending, attempting, and welding) on the corrosional properties on Ta	a magas
from an investigation of +3000 are presented, as a rolling, tube expanding, of brand TH3 and on Mb.	are data on the influence of technical operations (bending, attempting, and welding) on the corrosional properties on Ta	2

L 32996-66 EWT(1)/EWP(m) WW SOURCE CODE: UR/0170/66/010/005/0584/0591 ACC NRI AP60111985 AUTHOR: Leontivev. A. I.; Fedorov, V. K. ORG: Institute of Construction Physics, Moscow (Institut stroitelinoy fiziki, Moskva) TITLE: Experimental investigation of convective heat transfer in the movement of a gas in the inlet section of a cylindrical tube SOURCE: Inzhenerno-fizicheskiy zhurnal, v. 10, no. 5, 1966, 584-591 TOPIC TAGS: heat transfer, gas flow, thermodynamic analysis ABSTRACT: The experimental section consisted of a stainless steel tube with an outer diameter of 30 mm, an inside diameter of 24.3 mm, and a length of 1052 mm. A diagram of the equipment is given. All measurements were made under steady state conditions. Experiments were made at three values of the Mach number at the inlet of the tube: 0.28, 0.36, and 3. The temperature factor $T_{\rm cm}$ varied from 1 to 2.05 at $q_{\rm CT}={\rm const.}$ The following parameters were measured: p_0 , the stagnation pressure at the inlet of the tube; p_0 , the distribution of the static pressure along the length of the tube; $T_{\rm Cm}$, the temperature distribution at the wall over the length of the tube; $T_{\rm Cm}$, the stagnation temperature at the inlet 536.25 UDC: Card .1/2

。在大学经验的经济和经验的经验,但是在企业的经验的,但是是是一个人,我们就是是一个人,我们就是一个人,我们就是一个人,我们就是这个人,我们就是这个人,我们就是这个人,我们就是这个人,我们就是一个人,我们就是一个人,我们就

L 32996-66

ACC NR: AP6014985

of the tube; G, the sir flow rate; Q, the power supplied to the experimental section. The value of the specific heat flux was determined by the formula:

$$q_{ct} = (Q - Q')/\pi DI. \tag{1}$$

Initial conditions are shown in two large tables. According to the conclusion of the article, the general heat transfer law governing turbulent heat transfer in the boundary layer of the initial section

of a tube has the form:
$$St = \frac{0.014}{P_{es}^{0.55}} \left(\frac{1-\beta^{5}}{T_{er}}\right)^{0.5}. \quad (6)$$

Orig. art. has: 6 formulas, 3 figures and 2 tables.

003 SUBM DATE: 26Nov65/ ORIG REF: OTH REF: 011/ 20/ SUB CODE:

card 2/2 pla

CIA-RDP86-00513R000412630008-7" APPROVED FOR RELEASE: 03/20/2001

ACC NAL NIO	FNT(m)/EVIP(t)/E	SOVi	CE CODE: UR	/U413/66/000/o	Company T. 4	
······································	yayev, A.P.; Yukal	ov, I.N. Fede	rov, V.K.; Ys	khnina, V.D.;	Saparov, K.,	
Landa, A.F.					34	
					り事	
ORG: none		to No 1803	5 3	· .		
TITLE: None	agnetic iron. Cla	BE 40' NO TOO?	,		1066 61	
governor. Tea	breteniya, promys	hlennyye obraz	tsy, tovarnyy	e znaki, no.	, 1900, ta	
SOURCE: Izo	Interestrates &		llov	•		. 不動
TOPIC TAGS:	cast iron, nicke	T contaming a	Troj		nickel	
				on (in %):		1244
content. T	new nonmagnetic	OTTOWNIO -		3.0-3.1		0.004
				2.7-3.14		17.3
		M	anganese	6-8		
3		S		0.02-0.03	ī.	
			hosphorus 0 hromium	0.1-0.2		
			ickel	5-6	1	
			opper	2.0-2.5	Manual 1	100 1.25
				A 3-0-3/	/JPRS/	3 1 37 3
	11 / SUBM DATE:	3	lagnesiwi	0.1-0.14	C: 669.131.7	

			Belover V.K.	. Yekhnina. V	D.: Saparov	K.
. AUTHOR: Gu	lyayev, A.P.; Y	nkalov, I.N.	FOIDTOY VOICE	MeMye		. 1
Landa, A.F.						34
DRG: none				. 1		
TITLE: Nor	magnetic iron.	Class 40, No	180353			
	sobreteniya, pro			rnyye snaki, ne	. 7, 1966,	61 🔅
TOPIC TAGS	cast iron, ni	ckel containi	ng alloy			
A TO COMPA COMA	A new nonmagnet	ic cast iron	is proposed w	hich has a red	iced nickel	
ABSTRACLI		A the least of the	hemical compo	81610D (111 P)*		4.5
content.	A new nonmagnet This iron has th		Sautien.	2.0_2.1		
content.	rhis iron has th		Carbon Silicon	3.0-3.1 2.7-3.14		
content.	This iron has th		Silicon Manganese	2.7-3.14 6-8		
content.	this iron has th		Silicon Manganese Sulfur	2.7-3.14 6-8 0.02-0.03		
content.	This iron has th		Carbon Silicon Manganese Sulfur Phosphorus Chromium	2.7-3.14 6-8 0.02-0.03 0.05-0.66		
content.	this iron has th		Carbon Silicon Manganest Sulfur Phosphorus Chromium Nickel	2.7-3.14 6-8 0.02-0.03 0.05-0.06 0.1-0.2		
content.	this iron has the		Carbon Silicon Manganest Sulfur Phosphorus Chromium Nickel Copper	2.7-3.14 6-8 0.02-0.03 0.05-0.06 0.1-0.2		

FEDCROV, V. K.

Fedorov, V. K. - "The effect of average and large-sized doses of alcohol on the cortex of the brain," Trudy fiziol. laboratoriy im. Pavlova, Yol. XV, 1949, p. 194-228 Bibliog: 7 items

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949.)

FECORCY, Y. K. - "Investigation of the type of nervous activity in an unrestrained dog," Trudy fizibl. laboratoriy im. Pavlova, Vol. XV, 1949, p. 302-11, - Eibling: 6 items

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949.)

FEDOROV, V. K.

Fedorov, V. K. and Yakovleva, V. V. - "The analysis of the physiological mechanism of experimental neurosis," Trudy fiziol. laboratorii im. Pavlova, Vol. XV, 1949, p. 364-85, - Bibliog: p. 385

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949)

FEDOROV, V. K.

Fedorov, V. K. - "On the primary effect of narcotics (alcohol and chloral hydrate) on the large hemispheres of the brain," Trudy fiziol. laboratoriy im. Pavlofa, Vol. XV, 1949, p. 171-93, - Bibliog: 11 items

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949.)

Fractice, 7. K.

Fedorov, V. K. - "The relation of continued reflexes on the physical strength of positive irritants," Trudy fiziol. laboratoriy im. Pavlova, Vol. XV, 1949, p. 117-23

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949.)

FEDORO., V. K.

Fedorov, V. K. - "Conditional functional significance of the start, continuance and cessation of conditional irritants," Trudy fiziol. laboratoriy im. Pavlova, Vol. XV, 1949, p. 80-99, - Bibliog: 8 items

SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949.)

FEDUROV, V. K. - "Conditions relating to the appearance of internal retardation,"
Trudy fiziol. laboratoriy im. Pavlova, Vol. XV, 1949, p. 100-09, - Eibliog: 15 items
SO: U-4355, 14 August 53, (Letopis 'Zhurnal 'nykh Statey, No. 15, 1949.)


PEDOROV, V.K.

Effect of conditioned reflexes on the value of unconditioned salivary reflexes. Fixel. sh. SSSR 36 no.5:511-518 Sept-Oct 50 (CLML 20:4):

- 1. Physiological Department of the Institute of Experimental Medicine of the Academy of Medical Sciences, Leningrad.
- 2. Conditioned reflexes; relation between cortex and subcortical centers.

Method of studying conditioned reflex activity of mice., Zhur. vye, nerv, defat., 1, No. 5, 1951.

Monthly List of Russian Accessions, Library of Congress, April 1952.



FEDUROV, V. K.

"Achievements of the Past Twenty Years of Pharmacological Research in the Field of Higher Nervous activity." (Report No. 2, Pharmacology and Toxicology, Volume XIV, 1951, No. 3.

Extracted from Pharmacology of Ganglier Reflexes by S. V. Anichkov.

Fiziologicheskiy Zhurnal SSSR, Vol. 38, No. 1, pp 3-12, Jan/Feb 1952, Moscow/Leningrad

FEDOROV, V.K.

Investigation on the motility of nervous processes in mice. Fiziol. zh.SSSR 37 no.2:145-151 Mar-Apr 51. (CIML 21:1)

1. Laboratory of the Experimental Genetics of Higher Mervous Activity of the Institute of Physiology imeni I.P.Pavlov of the Academy of Sciences USSR.

FEDOROV. V.K.

On the training of responses of nervous processes in mice by multiple changes of a pair of reflexes. Fisiol.sh.SSSR 37 no.3:283-289 May-June 51. (CLML 21:1)

1. Institute of Physiology imeni I.P.Pavlov of the Academy of Sciences USSR, Leningrad.

LOMONOS, P.I.; FEDOROV, V.K.

Functional capacity of cortical cells in stimulation of larger hemispheres of the brain. Fixiol. sh. SSSR 37 no.5:579-586 Sept-Oct 51.

(CIML 21:4)

1. Physiology Department imeni I.P. Pavlov, Institute of Experimental Medicine, Academy of Medical Sciences USSR, Leningrad.

FEDOROV, V.K.

THE REST OF THE PERSON NAMED IN COLUMN TWO IS NOT THE PERSON NAMED IN COLUMN TWO IS NAMED IN THE PERSON NAMED IN COLUMN TWO IS NAM

Ansic principles of correlation of various motor reactions. Fisiol. sh. SSSR 38 no. 5:559-565 Sept-Oct 1952. (CIML 23:3)

1. Physiological Department imeni Academician I. P. Pavlov of the Institute of Experimental Sedicine, Academy of Medical Sciences USSR, Leningrad.

FEDOROV, V. K.

A STATE OF THE PARTY OF THE PAR

Problem of inheritance of acquired characteristics in mice.

Doklady Akad. nauk SSSR 84 no. 5:1061-1064 11 June 1952. (CLML 22:3)


1. Presented by Academician K. M. Bykov 16 April 1952. 2. Institute of Physiology imeni I. P. Pavlov, Academy of Sciences USSR.


FEDOROV. V.K.


Inheritance of changes in the higher nervous activity in animals. Izv.AB SSSR Ser.biol. no.5:3-15 S-0 '53. (MINA 6:9)

1. Institut fiziologii im. I.P.Pavlova.

(Nervous system) (Heredity)










Effect of exercising the nervous system of parents upon the mobility of neural processes in the descendants (mice). Trudy Inst.fixiol. (MIRA 7:5)

1. Laboratoriya eksperimental'noy genetiki vysshey nervnoy deyatel'nosti (zaveduyushchiy - V.K.Krasuskiy).

(Nervous system) (Inheritance of acquired characters)







GLEBOVSKIY, A.V.; FEDOROV, Vikt.K.

Comparative physiological investigations on the mobility of neural processes. Zhur. vys. nerv. deiat. 4 no.3:424-432 Hy-Je 154.

(MIRA 8:2)


I. Institut fiziologii imeni I.P.Pavlova Akademii nauk SSSR.

(NERVOUS SYSTEM, physiology.

lability of neural processes, determ. with conditioned reflex technic)

(REFLEX, CONDITIONED,

determ. of neural lability)



GLEBOVSKIY, A.V.; FEDOROV, Vikt. I. Method of comparative physiologic investigations of the higher nervous activity in animals. Zhur.vys.nerv.deiat. 4 no.5:581-585 JI-4g *54. 1. Institut fisiologii im. I.P.Pavlova Alcademii nauk SSSR. (CENTRAL MERVOUS SYSTEM, physiology, higher nervous funct. tests in animals)

PEDOROV, Vikt. K. Physiologic mechanism of the abatement of conditioned reflexes towards senility. Zmr. vys. nerv. defat. 4 no.4:568-573 Jl-Ag '54. (MIRA 8:3) 1. Institut fisiologii im. I.P.Pavlova Akademii nank SSSR. (AGING. age factor in conditioned reflex form. in mice) (REFLEX, COMDITIONED. age factor in reflex form. in mice)

Hereditary transmitting of modifications of the superior nervous activity in animals. Analele biol 9 no.2:55-69 Ap-Je 154.

MALYUGIHA, L.L.; MIRONOVA, A.I.; PEDODON, V.F.; SHARAD, L.M.

Significance of typologic characteristics of the higher nervous function in the formation and development of tumors produced by carcinogens in mice. Biul. eksp. biol. i med. 38 no.9:65-68 S 154.


(MIRA 7:12)

1. Is laboratorii eksperimental'noy genetiki vysshey nervnoy deyatel'nosti (sav. V.K.Krasuskiy) Instituta fiziologii imeni I.P.Pavlova (dir. akademik K.M.Bykov) AN SSSR i laboratorii eksperimental'noy onkologii (sav. chlen-korrespondent AME SSSR prof. L.M.Shabad) Instituta onkologii (dir. chlen-korrespondent AME SSSR prof. A.I. Serebrov) AME SSSR, Leningrad.

(NEOPLASMS, experimental,

higher nervous funct. in, role in form. & develop. of tumors)

(CHETRAL MERYOUS SYSTEM, function tests, typing of higher nervous funct., role in form & develop. of exper. tumors)



Name: FEDOROV, Viktor Konstantinovich

Dissertation: On the Physiology of the Higher Nervous Activity of Mice

Degree: Doc Biol Sci

Affiliation: /not indicated7

Defense Date, Place: 14 Oct 55, Council of the Inst of

Physiology imeni Pavlov, Acad Sci

Certification Date: 28 Apr 56

Source: BMVO 4/57

CIA-RDP86-00513R000412630008-7" APPROVED FOR RELEASE: 03/20/2001

SOFRONOV, M.S.; FEDOROV, Vict. K.

Effect of harmine on higher nervous activity in animals. Farm.
i teks. 18 m.3:3-9 My-Je '55. (MLEA 8:9)

1. Laboratoriya eksperimental'ney farmskelegii (zav. G.I. Tsobkalle) i laboratoriya eksperimental'ney genetiki vyschey nervnoy deyatel'nesti (zav. V.K. Krasuskiy) Institut fisiologii
I.P.Pavleva AM SSSR)

(RNFLEK, COMDITIONED,
off. of harmine)

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7

USSR/Medicine - Physiology FD-2710 Card 1/1 Pub. 33-19/28 Author : Glebovskiy, A. V,; Fedorov, Vikt. K. Title : A method for the study of the higher nervous activity of animals : Fiziol. zhur. 41, 104-108, Jan-Feb 1955 Periodical Abstract : Describes an electrodefensive method and apparatus for the study of conditioned reflex activity of animals such as mice, rats, guinea pigs, rabbits, cats, etc. Diagram; photograph; graphs. Institution : Institute of Physiology imeni I. P. Pavlov of the Academy of Sciences USSR Submitted : November 13, 1953

SOFROHOV, N.S.; FEDOROV, Viktor K. Hiffect of harmine on conditioned reflex activity in mice. Trudy Institute. 5:125-130 '56. (MIRA 10:1) 1. Leleratoriya eksperimental' noy farmakologii, maveduyushchiy - G.I.TSobkallo, i Leboratoriya eksperimental' noy genetiki vysshey nervnoy dayatel'nosti, maveduyushchiy - V.K.Krasuskiy (HARMINE) (GONDITIONED RESPONSE)

Effect of training the nervous system of mice on the higher nervous activity of their fourth-generation offspring. Trudy Instafiziol. 5:135-144 *56. (MERA 10:1) 1. Laboratoriya eksperimental noy genetiki vyashey nervnoy deyatel nosti. Zaveduyushchiy - V.K.Krasuskiy. (MERVOUS SYSTEM) (INHERITARCE OF ACQUIRED CHARGTERS)

"APPROVED FOR RELEASE: 03/20/2001

CIA-RDP86-00513R000412630008-7

USSR/Human and Initial Physiology. The Nervous System

T-12

Abs Jour : Ref Zhur - Riol., No 14, 1958, No 65723

Author

: Fedorov Vikt. K.

Inst

: :: :: USSR

Title

: Determination of the Type of Higher Nervous Activity in

Orig Pub : V sb.: Probl. fiziol. tsentr. mervn. sistemy. Moscow,

Loningrad, AN SSSR, 1957, 550-556

Abstract : Conditioned motor food reflexes were established in 4 mice to a light and sound stimulus lasting 45 seconds; theywere reinforced on the 30th second at two-minute intervals; discrimination was established and alteration performed, followed by errors. The experiments were conducted after one day's fast, the action time of the inhibitory signals was prolonged, the physical strength of the positive signals was increased, inhibitory or positive stimuli were reversed in the stereotype, and the interwals between them were reduced. In one mouse the norvous processes were strong,

Card

: 1/2

112

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7


USSR/Human and inimal Physiology. The Nervous System

T-12

Abs Jour : Ref Zhur - Riol., No 14, 1958, No 65723

balanced and active; in the second they were strong, unbalanced and active; in the third--strong, balanced and passive; and in the fourth-weak and passive.--M.I. Lisina

Card : 2/2



FEDOROV, V.K.

Comparative and physiological study of the higher nervous activity of animals. Trudy Vses. ob-va fiziol., biokhim. i farm. 4:99-104 158. (MIRA 14:2)

I. Institut fiziologii imeni I.P. Pavlova AN SSSR, Leningrad. (NERVOUS SYSTEM)

MALYUGINA, L.L., MIRONOVA, A.I., FEDOROV, Vikt. K. SHARAD, L.M.

Significance of typological characteristics of the higher nervous activity in the appearance and development of nouse mammary carcinoms. [with summary in English]. Biul.eksp.biol. i med. 45 no.6:85-89 Je '58 (MIEA 11:8)

1. In laboratorii eksperimental'noy genetiki vysshey nervncy deyatel'nosti (sav. V.K. Krasuskiy) Instituta fiziologii im. I.P. Psvlova
(dir. akad. K.M. Bykov) AM SSSR i laboratorii eksperimental'noy
onkologii (sav. - chlen-korrespondent AMM SSSR L.M. Shabad)
Instituta onkologii (dir. - chlen-korrespondent AMM SSSR A.I. Serebrov)
AMM SSSR, Leringrad. Predstavlena dystvitel'nym chelnom AMM SSSR
V.E. Chernigovskim.

(MEOPLASMS, experimental, mouse mammary carcinoma, eff. of type of higher nerv. activity (Rus))

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7

AUTHORS: Nemchinova, I. I., Pedorov, Vikt. K. SOV/20-121-1-48/55

TITLE: On the Problem of Sex Relation Among the Progeny of Mice Endowed With Different Functional Properties in Their Nervous Systems

(E voprosu o sootnoshenii polov v potomstve u myshey s raz-

lichnymi funktaonal'nymi svoystvami nervnoy sistemy)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol. 121, Nr 1,

pp. 169 - 171 (USSR)

The physiological differences between the male and female individuals of either sex have been investigated in many papers. In recent time for the first time investigations were carried out which are to clarify the influence of the nervous system

on the sex relation of the progeny. Thus, it was proved (Refs 3,4) that this relation may be changed conditional-reflexly within considerable limits. Unfortunately, this communication was

confirmed neither by the author himself nor by other researchers.

In the present paper the authors give facts on this. The second author paid special attention to the problem of the mobility of the nervous processes in his many years' investigations of the higher nervous activity of rodents (Ref 6).

Card 1/4

ABSTRACT:

On the Problem of Sex Relation Among the Progeny of Mice SOV/20-121-1-48/55 Endowed With Different Functional Properties in Their Nervous Systems

This mobility anables the animal to adapt itself perfectly to the environmental conditions. According to I.P.Pavlov this property of the nervous system is characterized by the velocity of the substitution of the cortical processes of the nerves-stimulation and inhibition - and may be determined by means of various methods. One of these methods is the mutual reformation of the conditional reflexes. Mice of different oncological lines were subjected to the investigation; A and C.HA of high sensitivity to cancer which furnish a rather high percentage of spontaneous ulcers (Refs 2,5), and C57 which are less sensitive to cancer and homozygote with respect of the lacking of the "milk factor". It was proved earlier (Ref 7) that the mobility of the nervous processes of females of a high sensitivity to cancer is greater than of females which are less sensitive to cancer. The mean velocity of the reformation of the conditional reflexes in the case of 91 females of the line A amounts to 2,7 \pm 2,19 experiments, in the case of 57 females of the line C_2HA to 19,5 \pm 0,91 experiments, and in the case of 163 females of the line 1056

Card 2/4

On the Problem of Sex Relation Among the Progeny of Mice SOV/20-121-1-48/55 Endowed With Different Functional Properties in Their Nervous Systems

to only 15,4 ± 0,86 experiments. With reference to these obvious differences between the lines of different sensitivity to cancer the authors counted how many male and female descendents existed in order to clarifythe connection between the degree of mobility and the sex relation of the progeny (Table 1). On the strength of this material it was assumed that the females with higher mobility produce more males, whereas the more inert parents produce more females. This was confirmed statistically by further investigations. There are 2 tables and 7 Soviet references.

ASSOCIATION: Institut fiziologii im. I.P.Pavlova Akademii nauk SSSR

(Institute of Physiology imeni I.P.Pavlov, AS USSR)

PRESENTED: March 28, 1958, by K.M.Bykov, Member, Academy of Sciences,

USSR

SUBMITTED: March 8, 1958

Card 3/4

"APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7

On the Problem of Sex Relation Among the Progeny of Mice SOV/2o-121-1-48/55 Endowed With Different Functional Properties in Their Nervous Systems

1. Sex--Genetic factors
--Physiological effects
4. Nervous system--Physiology
5. Cancer--Physiological


Card 4/4

FEDOROV, VIKE.K. Effect of single and continuous application of small and medium doses of caffein on the extinctive inhibition in mice. Nauch. (MIRA 14:10)

soob. Inst. fixiol. AN SSSR no.1:72-74 159.

1. Gruppa po izucheniyu senetiki vysshey nervnoy deyatel'nosti u gryzunov (zav: Vikt.K.Fedorov)Instituta fiziologii imeni Pavlova AN SSSR. (CONDITIONED RESPONSE) (CAFFEIN-PHYSIOLOGICAL EFFECT)

CIA-RDP86-00513R000412630008-7" APPROVED FOR RELEASE: 03/20/2001



17(1) AUTHORS: Ol'nyanskaya, R. P., Fedorov, Vikt. K. SOV/20-124-1-68/69 TITLE Basal Metabolic Rate and Typological Characteristics of the Rervous System in Mice (Osnovnoy obmen i tipologicheskiye osobennosti nervnoy sistemy u myshey) PERIODICAL: Doklady Akademii nauk SSSR, 1959, Vol 124, Nr 1, pp 237-240 (USSR) ABSTRACT: It is the aim of the present paper to detect which of the characteristic features mentioned in the title find particular expression in the reflex reactions of the respiratory change. The experiment was carried out with 29 male mice of the line C57. To begin with, the conditionally reflex activity of the animals was investigated according to the "motor alimentary method" (dvigatel'no-pishchevaya metodika)(Refs 1, 2, 7). During 2 - 2.5 months the respiratory change was observed in climate chambers for one hour every day. During this procedure the animals were in quiescent state. As can be seen from table one all animals were divided into two groups, according to the results of respiratory change: I. Nice whose basal metabolic Card 1/3 rate reached a certain level during the first 18.2 days

APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7"

Basal Metabolic Rate and Typological Characteristics 507/20-124-1-68/69 of the Nervous System in Mice

(the average duration until the basal metabolic rate level becomes stable). Low level and low variability coefficient were characteristic of this group. II. For this group of animals no stable level of the basal metabolic rate was achieved. From the results obtained can be concluded that the typological characteristics of the nervous system can be seen from the reactions of the respiratory change. Those characteristic features are particularly expressed by the time the basal metabolic rate requires to attain a stable level. The rate of development of the positive "motor alimentary reflex" and the time required by the basal metabolic rate to attain a stable level coincide most considerably. A less important correlative relation was observed between the rate of change of the conditional reflexes and the mentioned adjustment of the level. The final solution of the problem is left to further research. The results obtained are a first attempt towards revealing the

Card 2/3

Basal Metabolic Rate and Typological Characteristics SOV/20-124-1-68/69 of the Rervous System in Mice

relation between the individual properties of the nervous system and the processes of respiratory change. There are 1 figure, 2 tables, and 7 Soviet references.

ASSOCIATION: Institut fimiologii im. I. P. Pavlova Akademii nauk SSSR (Institute of Physiology imeni I. P. Pavlov, Academy of Sciences, USSR)

PRESENTED: September 8, 1958, by K. M. Bykov, Academician

SUPMITTED: August 20, 1958

Card 3/3

Some problems with regard to the present state of studies on the type of the nervous system in animals. Fixiol. zhur. [Ukr.] 6 no. 5:571-576 S-0 160. (MIRA 13:10)

1. Institut fiziologii im. I.P. Pavlova Akademii nauk SSSR, Leningrad.

(MERVOUS SYSTEM)

Augis 18

The same state of the same of

FEDOROV, Vikt.K.

Comparison of the results of individual studies in the evaluation of the basic properties of the higher nervous activity in mice.

Zhur. vys. nerv. deiat. 11 no.4:746-752 J1-Az '61. (MLM 15:2)

1. Pavlov Institute of Physiology, U.S.S.R. Academy of Sciences, Koltushi. (MERYOUS SYSTEM) (CONDITIONED MESTONEE)

APPROVED FOR RELEASE: 03/20/2001 CIA-RDP86-00513R000412630008-7"

राज्यकाका प्रमानेत कारक साम निर्माणका स्थापका वृत्ताका व प्रमानिक प्रतास ।

PEDOROV, Vikt.K.

Study of higher nercous activity in the offspring of trained and standard parents during four generations by the method of motor defense conditioned reflexes. Trudy Inst. fiziol. 10:143-151'62 (MIRA 17:3)

1. Gruppa fiziologii tipa vysshey nervnoy deyatel nosti zhivotnykh (zav. - Vikt.K.Fedorov) Instituta fiziologii imeni Pavlova AN SSSR.

Study of typological properties of the nervous system and gas exchange in rodents. Trudy Inst. fisiol. 10:255-264 162 (MIRA 17:3)

1. Laboratoriya nevrofisiologicheskikh problem (zav. - K.M. Bykov [deceased]) i gruppa po izucheniyu genetiki vysshey nervnoy deyatel nosti i gryzumov (zav. - Vikt.K.Fedorov) Instituta fisiologii imeni Pavlova AN SSSR.

FEDOROV, V. K.

Appropos of V.L. Merkuloy's article, "Materials on I.P.
Pavlov's attitude towards the use of mathematical methods
in studying problems in digestion and higher nervous activity."
Zhur. nevr. 1 psikh. 62 no.3:478-479 '62. (MIRA 15:3)
(NERVOUS SYSTEM) (DIGESTION) (MATHEMATICS)
("ERKULOV, V.L.)

